

Temperature and latitude analysis to predict potential spread and seasonality for COVID-19

Mohammad M. Sajadi, MD,^{1,2} Parham Habibzadeh, MD,³ Augustin Vintzileos, PhD,⁴ Shervin Shokouhi, MD,⁵ Fernando Miralles-Wilhelm, PhD,^{6,7} Anthony Amoroso, MD^{1,2}

¹ Institute of Human Virology, University of Maryland School of Medicine, Baltimore, USA

² Global Virus Network (GVN), Baltimore, USA

³ Persian BayanGene Research and Training Center, Shiraz University of Medical Sciences, Shiraz, Iran

⁴ Earth System Science Interdisciplinary Center, University of Maryland, College Park, USA

⁵ Infectious Diseases and Tropical Medicine Research, Shaheed Beheshti University of Medical Sciences, Tehran, Iran

⁶ Department of Atmospheric and Oceanic Science, University of Maryland, College Park, USA

⁷ The Nature Conservancy, Arlington, USA

Corresponding author:

Mohammad Sajadi, MD
Associate Professor
Institute of Human Virology
Global Virus Network (GVN) Center of Excellence
University of Maryland School of Medicine
725 W. Lombard St. (N548)
Baltimore, MD 21201
Office (410) 706-1779
Fax (410) 706-1992
msajadi@ihv.umaryland.edu

Abstract

A significant number of infectious diseases display seasonal patterns in their incidence, including human coronaviruses. We hypothesize that SARS-CoV-2 does as well. To date, Coronavirus Disease 2019 (COVID-19), caused by SARS-CoV-2, has established significant community spread in cities and regions only along a narrow east west distribution roughly along the 30-50 N” corridor at consistently similar weather patterns (5-11°C and low specific and absolute humidity). There has been a lack of significant community establishment in expected locations that are based only on population proximity and extensive population interaction through travel. We have proposed a simplified model that shows a zone at increased risk for COVID-19 spread. Using weather modeling, it may be possible to predict the regions most likely to be at higher risk of significant community spread of COVID-19 in the upcoming weeks, allowing for concentration of public health efforts on surveillance and containment.

Background:

Many infectious diseases show a seasonal pattern in their incidence. An onerous burden for health care systems around the globe, influenza is the characteristic example.¹ The influenza virus shows significant seasonal fluctuation in temperate regions of the world but nevertheless displays less seasonality in tropical areas.²⁻⁴ Despite the multitude of possible mechanisms proposed to explain this variation, our current understanding of this phenomenon is still superficial.⁵

Coronavirus Disease 2019 (COVID-19), caused by SARS-CoV-2, initially came to attention in a series of patients with pneumonia of unknown etiology in the Hubei province of China, and subsequently spread to many other regions in the world through global travel.⁶ Because of geographical proximity and significant travel connections, epidemiological modeling of the epicenter predicted that regions in Southeast Asia, and specifically Bangkok would follow Wuhan, and China in the epidemic.⁷ More recently, the World Health Organization has declared this as a pandemic. (WHO, 2020) For many the biggest concern is not how large the problem but what will happen in the coming months and which areas and populations are most at risk.

A number of studies, both laboratory,⁸ epidemiological studies,⁹ and mathematical modelling,¹⁰ point to role of ambient temperature on the survival and transmission of viruses. The tremendous level of research supporting both ambient temperature and humidity in its role in transmission and infection motivated this study to examine the influence of environmental factors on COVID-19. We sought to determine whether climate could be a factor in the spread of this disease.

Methods:

2-meter (2m) temperatures, relative humidity (RH), specific humidity (Q), and absolute humidity (AH) were based on data from the ECMWF ERA-5 reanalysis. Climatologic (1979-2019) and persistence forecasting (2019 data) was used to analyze latitude and temperature trends globally and for affected areas using ERA-3 (Interim) and ERA-5. ERA-3 reanalysis data obtained from Climate Reanalyzer (<https://ClimateReanalyzer.org>), Climate Change Institute, University of Maine, USA. ERA-Interim reanalysis data (<https://doi.org/10.1002/qj.828>) cover the earth with a resolution of 80 km x 80 km. The analysis of 2-meter temperature is performed in separate analysis following the upper air 4D-Var analysis. ERA-5 reanalysis data (C3S, 2017) covers the earth with a resolution of 30 km x 30 km. Preliminary daily updates are available 5 days of real time though quality-assured monthly updates are published within 3 months of real time. 2m temperature was calculated by interpolating between the lowest model level and the Earth's surface, taking into account the atmospheric conditions.

2-meter Temperature (2m) is Temperature at the height of 2 meters, 1000 hPa Temperature is the temperature at the pressure level of 1000 hPa and is calculated by interpolation from the numerical model levels. Relative humidity (RH) is the percentage of the maximum amount of water vapor that the atmosphere can hold at a given temperature (saturation). Specific humidity (Q) is defined as the mass of water vapour in a unit mass of moist air (g/kg). Absolute humidity (AH) is defined as the total mass of water vapor present in a given volume or mass of air (g/m³). Significant community transmission is defined as >6 reported death as of March 5, 2020.

Results:

Through March 5, 2020, significant community transmission has occurred in a consistent east and west pattern. The new epicenters of disease were all roughly along the 30-50° N zone; to South Korea, Japan, Iran, and Northern Italy (Figure 1).¹¹ After the unexpected emergence of a large outbreak in Iran, we first made this map in late February. Since then new areas with significant community transmission include the Northwestern United States and France (Figure 1). Notably, during the same time, COVID-19 failed to spread significantly to countries immediately north and south of China. The number of patients and reported deaths in Southeast Asia is much less when compared to more temperate regions noted above.¹¹

1000hPa Temperature (°C)
NDJFM 2019

ECMWF ERA-Interim




Figure 1. World temperature map November 2018-March 2019. Color gradient indicates 1000hPa temperatures in degrees Celsius. Black circles represent countries with significant community transmission (> 6 deaths as of 3/5/2019). Image from Climate Reanalyzer (<https://ClimateReanalyzer.org>), Climate Change Institute, University of Maine, USA.

Further analysis using 2-meter (2m) temperatures from 2020 rather than hPa temperatures yields similar results (Figure 2). In the months of January 2020 in Wuhan and February 2020 in the other affected, there is a striking similarity in the measures of average temperature (4-9 °C at the airport weather stations, and with city temperatures slightly higher due to urban effect,¹² they are within a range of 5-11 °C) and low specific and absolute humidity (Table 1). In addition to having similar average temperature, humidity, and latitude profiles, these locations also exhibit a commonality in that

the timing of the outbreak coincides with a nadir in the yearly temperature cycle, and thus with relatively stable temperatures over a more than a one month period of time (Table 1).

Average 2-meter Temperature (°Celsius) for Jan-Feb 2020 (ERA-5)




Figure 2. World temperature map January 2020-February 2020. Color gradient indicates 2-meter temperatures in degrees Celsius based on data from the ECMWF ERA-5 reanalysis. White circles represent countries with significant community transmission (> 6 deaths as of 3/5/2020), and red isolines areas with temperature between 5-11°C. Generated using Copernicus Climate Change Service Information 2020.

Given the temporal spread among areas with similar temperature and latitude, some predictions can tentatively be made about the potential community spread of COVID-19 in the coming weeks. Using 2019 temperature data for March and April, risk of community spread could be predicted to affect areas just north of the current areas at risk (Figure 3). These could include (from East to West) Manchuria, Central Asia, the Caucasus, Eastern Europe, Central Europe, the British Isles, the Northeastern and Midwestern United States, and British Columbia. However, this simplified analysis does not take into account the effect of warming temperatures. The marked drop in cases in Wuhan could well be linked to corresponding recent rising temperatures there (Table 1).




Figure 3. World 2 meter average temperature map March 2019-April 2019 predicting at risk zone for March-April 2020. Color gradient indicates average 2M temperatures in degrees Celsius. Tentative zone at risk for significant community spread in the near-term include land areas within the green bands, outlined in dark black (showing 5-11°C zone based on 2019 data), and will change based on actual average temperatures during this time period. Image from Climate Reanalyzer (<https://ClimateReanalyzer.org>), Climate Change Institute, University of Maine, USA. Digital manipulation by Cameron Gutierrez and Glenn Jameson.

City	Nov 2019				Dec 2019				Jan 2020				Feb 2020			
	2M (°C)	Rh (%)	Q (g/kg)	AH (g/m³)	2M (°C)	Rh (%)	Q (g/kg)	AH (g/m³)	2M (°C)	Rh (%)	Q (g/kg)	AH (g/m³)	2M (°C)	Rh (%)	Q (g/kg)	AH (g/m³)
Wuhan	14	66	6	8	8	74	5	6	5	84	4	6	10	77	6	7
Tokyo	14	72	7	9	10	73	5	7	8	72	5	6	9	66	5	6
Qom	9	61	5	5	7	72	5	6	4	69	4	4	8	44	4	4
Milan	9	85	6	8	7	80	5	6	5	77	4	5	8	60	4	5
Daegu	9	68	5	6	2	62	3	4	3	67	3	4	4	62	3	4
Seattle	7	84	5	7	7	88	5	7	7	85	5	7	6	82	5	6
Mulhouse	7	89	5	6	7	87	5	6	2	87	4	5	6	73	5	6

Large cities tentatively predicted to be at risk in the coming weeks

London	7	90	6	7	7	89	5	7	7	89	6	7	7	81	5	6
Manchester	6	92	5	7	5	91	5	6	6	90	5	7	6	84	5	6
Berlin	6	88	5	7	4	84	4	6	4	84	4	5	6	77	5	6
Prague	6	86	5	6	3	82	4	5	2	84	4	5	5	72	4	5
Hamburg	6	91	5	7	5	87	5	6	5	89	5	6	6	82	5	6
Vancouver	8	81	5	7	6	85	5	6	5	82	5	6	5	81	4	6
New York	7	65	4	5	3	74	4	5	3	69	4	4	4	69	4	5
Warsaw	6	88	5	7	3	86	4	5	2	87	4	5	4	79	4	5
Glasgow	5	88	5	6	6	89	5	7	7	86	5	7	5	85	5	6
Kiev	5	84	5	6	3	87	4	5	1	86	3	4	2	77	4	4
St. Louis	5	71	4	5	3	75	4	5	2	78	4	4	2	72	3	4
Beijing	5	53	3	4	-3	54	2	2	-3	58	2	2	1	62	2	3

Previously predicted city where COVID-19 failed to take hold

Bangkok	28	70	16	19	26	70	15	17	28	74	17	20	28	70	16	19
---------	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----

Table 1. November 2019 to February 2020 average 2m temperature (°C), relative humidity (RH, %), Specific humidity (Q, g/kg), and absolute humidity (AH, g/m³) data from cities with community spreading of COVID-19 and those at potentially at risk (as of 3/5/20). Temperature and humidity based on data from the ECMWF ERA-5 reanalysis.

Discussion: The association between temperature in the cities affected with COVID-19 deserves special attention. There is a similarity in the measures of average temperature (5-11°C) and RH (47-79%) in the affected cities and known laboratory conditions that are conducive to coronavirus survival (4°C and 20-80% RH).¹³ In the time we have written up these results, new centers of significant community outbreak include parts of Northeastern United States, Spain, Germany, and England, all of which had seen average temperatures between 5-11°C in January and February 2020, and included in either the Jan-Feb 2020 map (Figure 2), or Mar-Apr risk map (Figure 3).

Temperature and humidity are also known factors in SARS-CoV, MERS-CoV and influenza survival.¹⁴⁻¹⁷ Furthermore, new outbreaks occurred during periods of prolonged time at these temperatures, perhaps pointing to increased risk of outbreaks with prolonged conditions in this range. Besides potentially prolonging half-life and viability of the virus, other potential mechanisms associated with cold temperature and low humidity include stabilization of the droplet and enhanced propagation in nasal mucosa, as has been demonstrated with other respiratory viruses.^{8,18} Finally, even colder areas in the more Northern latitudes have been relatively free of COVID-19 pointing to a potential minimum range for the temperature, which could be due to avoidance of freeze-thaw cycles that could affect virus viability or other factors (at least one human coronavirus tested is freeze-thaw resistant).¹⁹ All of the above points to a potential direct relation between temperature and SARS-CoV-2 environmental survival and spreading. This hypothesis can be tested in experimental conditions similar to work that has been done before,¹³ environmental sample testing from areas of ongoing infection, and close epidemiologic and climate studies in the coming weeks.

In the coming 2 months, temperatures will rise dramatically across many areas in the Northern Hemisphere. However, areas to the north which develop temperature profiles that may now overlap the current areas at risk only transiently as they rapidly warm (with possible exception of areas such as the Northwest United States and British Columbia, which can show prolonged cyclical nadirs). Furthermore, as the virus moves further north it will encounter sequentially less human population densities. The above factors, climate variables not considered or analyzed (cloud cover, maximum temperature, etc.), human factors not considered or analyzed (impact of epidemiologic interventions, concentrated outbreaks like cruise ships, travel, etc.), viral factors not considered or analyzed (mutation rate, pathogenesis, etc.), mean that although the current correlations with latitude and temperature seem strong, a direct causation has not been proven and predictions in the near term are speculative and have to be considered with extreme caution.

Human coronaviruses (HCoV-229E, HCoV-HKU1, HCoV-NL63, and HCoV-OC43), which usually cause common cold symptoms, have been shown to display strong winter seasonality between December and April, and are undetectable in summer months in temperate regions.²⁰ Some studies have shown that the alphacoronavirus HCoV-229E peaks in the fall, while HCoV-OC43 (a betacoronavirus in the same genera as SARS-CoV-2) has a winter predominance.^{21,22} Although it would be even more difficult to make a long-term prediction at this stage, it is tempting to expect COVID-19 to diminish considerably in affected areas (above the 30° N") in the coming months and into the summer. It could perhaps prevail at low levels in tropical regions similar to influenza, cause outbreaks in the Southern Hemisphere at the same time, and begin to rise again in late fall and winter in temperate regions in the upcoming year. One other possibility is that it will not be able to sustain itself in the summer in the tropics and Southern Hemisphere and disappear, just as SARS failed to do so in 2003; however, the ever-increasing number of cases worldwide make this increasingly less likely. MERS has pointed to as a case of a betacoronavirus that is able to spread in all seasons. However, it should be remembered that the vast majority of cases were in the Arabian Peninsula, which has a Hot desert climate (Köppen classification), and that influenza infection there does not follow the same pattern as more temperate climates.²³ In the upcoming summer months in the Northern Hemisphere, surveillance efforts for

SARS-CoV-2 in currently affected areas will be important to determine if there is a viral reservoir (such as prolonged stool shedding). Similarly, surveillance efforts in the tropics, as well as New Zealand, Australia, South Africa, Argentina, and Chile between the months of June through September may be of value in determining establishment in the human population.

An avenue for further research involves the use of integrated or coupled epidemiological-earth-human systems models, which can incorporate climate and weather processes and variables (e.g., dynamics of temperature, humidity) and their spatiotemporal changes, as well as simulate scenarios of human interactions (e.g., travel, transmission due to population density). Such models can assimilate data currently being collected to accelerate the improvements of model predictions on short time scales (i.e., daily to seasonal). This type of predictive approach would allow to explore questions such as what are population centers most at risk and for how long; where to intensify large scale surveillance and tighten control measures to prevent spreading; better understanding of limiting factors for virus spreading in the southern hemisphere; and making predictions for a 2021-2022 virus season. A better understanding of the cause of seasonality for coronaviruses and other respiratory viruses would undoubtedly aid in better treatments and/or prevention, and be useful in determining which areas need heightened surveillance.

Conflict of interest: None to declare.

M.M.S supported by NIH grant 1R01AI147870-01A1.

Acknowledgements: Digital manipulation of Figure 3 courtesy of Cameron Gutierrez and Glenn Jameson. The authors would also like to thank Betsy Weatherhead (Senior Scientist - retired, U. Colorado at Boulder) for analytical assistance on examining global COVID-19 and putting data in the context of environmental factors.

References

1. Collaborators GBDI. Mortality, morbidity, and hospitalisations due to influenza lower respiratory tract infections, 2017: an analysis for the Global Burden of Disease Study 2017. *Lancet Respir Med* 2019; **7**(1): 69-89.
2. Viboud C, Alonso WJ, Simonsen L. Influenza in tropical regions. *PLoS Med* 2006; **3**(4): e89.
3. Bloom-Feshbach K, Alonso WJ, Charu V, et al. Latitudinal variations in seasonal activity of influenza and respiratory syncytial virus (RSV): a global comparative review. *PLoS One* 2013; **8**(2): e54445.
4. Li Y, Reeves RM, Wang X, et al. Global patterns in monthly activity of influenza virus, respiratory syncytial virus, parainfluenza virus, and metapneumovirus: a systematic analysis. *Lancet Glob Health* 2019; **7**(8): e1031-e45.
5. Tamerius J, Nelson MI, Zhou SZ, Viboud C, Miller MA, Alonso WJ. Global influenza seasonality: reconciling patterns across temperate and tropical regions. *Environ Health Perspect* 2011; **119**(4): 439-45.
6. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; **395**(10223): 497-506.
7. Bogoch, II, Watts A, Thomas-Bachli A, Huber C, Kraemer MUG, Khan K. Potential for global spread of a novel coronavirus from China. *J Travel Med* 2020.
8. Lowen AC, Mubareka S, Steel J, Palese P. Influenza virus transmission is dependent on relative humidity and temperature. *PLoS Pathog* 2007; **3**(10): 1470-6.
9. Barreca AI, Shimshack JP. Absolute humidity, temperature, and influenza mortality: 30 years of county-level evidence from the United States. *Am J Epidemiol* 2012; **176** Suppl 7: S114-22.
10. Zuk T, Rakowski F, Radomski JP. Probabilistic model of influenza virus transmissibility at various temperature and humidity conditions. *Comput Biol Chem* 2009; **33**(4): 339-43.

11. Coronavirus COVID-19 Global Cases by Johns Hopkins CSSE. 2020.
<https://gisanddata.maps.arcgis.com/apps/opsdashboard/index.html#/bda7594740fd40299423467b48e9ecf6> (accessed 3/3/2020).
12. Aude Lemonsu MD, Deque M, Somot S, Alias A, Masson V. Benefits of explicit urban parameterization in regional climate modeling to study climate and city interactions. *Climate Dynamics* 2018; **52**: 2745-64.
13. Casanova LM, Jeon S, Rutala WA, Weber DJ, Sobsey MD. Effects of air temperature and relative humidity on coronavirus survival on surfaces. *Appl Environ Microbiol* 2010; **76**(9): 2712-7.
14. Otter JA, Donskey C, Yezli S, Douthwaite S, Goldenberg SD, Weber DJ. Transmission of SARS and MERS coronaviruses and influenza virus in healthcare settings: the possible role of dry surface contamination. *J Hosp Infect* 2016; **92**(3): 235-50.
15. Chan KH, Peiris JS, Lam SY, Poon LL, Yuen KY, Seto WH. The Effects of Temperature and Relative Humidity on the Viability of the SARS Coronavirus. *Adv Virol* 2011; **2011**: 734690.
16. van Doremalen N, Bushmaker T, Munster VJ. Stability of Middle East respiratory syndrome coronavirus (MERS-CoV) under different environmental conditions. *Euro Surveill* 2013; **18**(38).
17. Tan J, Mu L, Huang J, Yu S, Chen B, Yin J. An initial investigation of the association between the SARS outbreak and weather: with the view of the environmental temperature and its variation. *J Epidemiol Community Health* 2005; **59**(3): 186-92.
18. Schaffer FL, Soergel ME, Straube DC. Survival of airborne influenza virus: effects of propagating host, relative humidity, and composition of spray fluids. *Arch Virol* 1976; **51**(4): 263-73.
19. Lamarre A, Talbot PJ. Effect of pH and temperature on the infectivity of human coronavirus 229E. *Can J Microbiol* 1989; **35**(10): 972-4.
20. Gaunt ER, Hardie A, Claas EC, Simmonds P, Templeton KE. Epidemiology and clinical presentations of the four human coronaviruses 229E, HKU1, NL63, and OC43 detected over 3 years using a novel multiplex real-time PCR method. *J Clin Microbiol* 2010; **48**(8): 2940-7.
21. Vabret A, Mourez T, Gouarin S, Petitjean J, Freymuth F. An outbreak of coronavirus OC43 respiratory infection in Normandy, France. *Clin Infect Dis* 2003; **36**(8): 985-9.
22. Cabeça TK, Granato C, Bellei N. Epidemiological and clinical features of human coronavirus infections among different subsets of patients. *Influenza Other Respir Viruses* 2013; **7**(6): 1040-7.
23. Caini S, El-Guerche Seblain C, Ciblak MA, Paget J. Epidemiology of seasonal influenza in the Middle East and North Africa regions, 2010-2016: Circulating influenza A and B viruses and spatial timing of epidemics. *Influenza Other Respir Viruses* 2018; **12**(3): 344-52.